

RESEARCH ARTICLE

Relationship between fall history and toe grip strength in older adults with knee osteoarthritis in Japan: A cross-sectional study

Yuya Mawarikado¹, Yusuke Inagaki^{2*}, Tadashi Fujii³, Takanari Kubo⁴, Akira Kido², Yasuhito Tanaka^{1,5}

1 Graduate School of Medicine, Musculoskeletal Reconstructive Surgery, Nara Medical University, Nara, Japan, **2** Department of Rehabilitation Medicine, Nara Medical University, Nara, Japan, **3** Department of Orthopaedic Surgery, Kashiba Asahigaoka Hospital, Nara, Japan, **4** Department of Rehabilitation, Osaka Kawasaki Rehabilitation University, Osaka, Japan, **5** Department of Orthopaedic Surgery, Nara Medical University, Nara, Japan

* yinagaki@naramed-u.ac.jp



Abstract

Background

Knee osteoarthritis (KOA), one of the most common musculoskeletal diseases in older adults, is associated with a high incidence of falls. Similarly, toe grip strength (TGS) is associated with a history of falls in older adults; however, the relationship between TGS and falls in older adults with KOA who are at risk of falling is not known. Therefore, this study aimed to determine if TGS is associated with a history of falls in older adults with KOA.

Methods

The study participants, older adults with KOA scheduled to undergo unilateral total knee arthroplasty (TKA), were divided into two groups: non-fall (n = 256) and fall groups (n = 74). Descriptive data, fall-related assessments, modified Fall Efficacy Scale (mFES), radiographic data, pain, and physical function including TGS were evaluated. The assessment was conducted on the day before performing TKA. Mann–Whitney and chi-squared tests were performed to compare the two groups. Multiple logistic regression analysis was performed to determine the association of each outcome with the presence or absence of falls.

Results

Mann–Whitney U test revealed that the fall group had statistically significantly lower height, TGS on the affected and unaffected sides, and mFES. Multiple logistic regression analysis revealed that the incidence of fall history is associated with TGS on the affected side; the weaker the affected TGS of the KOA, the more likely the individual is to fall.

OPEN ACCESS

Citation: Mawarikado Y, Inagaki Y, Fujii T, Kubo T, Kido A, Tanaka Y (2023) Relationship between fall history and toe grip strength in older adults with knee osteoarthritis in Japan: A cross-sectional study. PLoS ONE 18(3): e0282944. <https://doi.org/10.1371/journal.pone.0282944>

Editor: Faizan Iqbal, Baqai Medical University, PAKISTAN

Received: November 1, 2022

Accepted: February 27, 2023

Published: March 13, 2023

Copyright: © 2023 Mawarikado et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its [Supporting Information](#) files.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Conclusions

Our results indicate that TGS on the affected side is related to a history of falls in older adults with KOA. The significance of evaluating TGS among patients with KOA in routine clinical practice was demonstrated.

Introduction

Falls in older adults can lead to injuries, such as bone fractures, and to significant deterioration of physical function. Over 30% of people aged 65 or older experience approximately one fall every year [1]. Knee osteoarthritis (KOA) is one of the most common musculoskeletal diseases in older adults. It has been reported that an increased history of falls and hip fractures is associated with an increase in knee pain, and the incidence of non-vertebral fractures is 1.6 times higher in older adults with KOA than in those without [2]. The incidence of falls in older adults with KOA is approximately 30% higher than that in healthy older adults, and approximately half of individuals aged 60 years or older fall at least once annually [3, 4]. Older adults with KOA and a fall history are more likely to fall after total knee arthroplasty (TKA) than a cohort of older adults with KOA who do not have a history of falls [5, 6].

As mentioned, older adults with KOA have a high incidence of falls; however, related risk factors remain undetermined in this population. Knee pain, impaired balance, lower muscle weakness, and decreased walking ability have been reported as risk factors for falls in individuals with KOA [7, 8]. Previous studies on older adults without KOA indicated that clinical benchmarks, such as Timed Up and Go (TUG) [9, 10], fall-related self-efficacy [11], knee extension strength [12], and toe grip strength (TGS) [13–15] are associated with the risk of a fall. These outcomes may be even stronger predictors in individuals with KOA, who generally have impaired mobility and muscle strength.

Among the aforementioned risk factors for falls, particular attention has been paid to TGS. Many studies reported an association between TGS and falls in older adults. With increasing age, TGS in older adults deteriorates [16–18], resulting in decreased walking speed and static balance ability [19–21]. TGS training to nursing home residents led to significant improvements in the fall risk index (-1.4 points). A significant increase in TGS was noted in the intervention group compared to that in the non-intervention group (Intervention group increased 1.9 kg, non-intervention group decreased 0.2 points) [22]. Therefore, it is clinically important to evaluate and strengthen TGS to prevent falls in older adults.

To date, no study has investigated the relationship between falls and TGS in older adults with KOA who are at risk of falling owing to lower limb muscle weakness and impaired balance. We hypothesized that the history and frequency of falls in older adults with KOA would be associated with TGS. Therefore, this study aimed to determine whether TGS is associated with a history of falls in older adults with KOA.

Methods

Ethical issues

This study complied with the Declaration of Helsinki and was approved by the Research Ethics Committee of Kashiba Asahigaoka hospital (2019-04-21-007). Details of the study protocol and aim were explained to all participants, both verbally and in writing. All study participants then signed a written consent prior to participating in the study.

Participants

This study used a descriptive cross-sectional design to identify the association between falls and clinical evaluation factors including TGS. We recruited 407 participants with KOA, scheduled to undergo unilateral TKA at a single hospital in Japan between May 2019 and September 2021. The inclusion criteria were: 1. diagnosis of KOA, 2. ability to ambulate independently or with a T-cane at the time of pre-operative evaluation, 3. individuals who were scheduled for primary TKA, 4. individuals between 60 and 84 years of age, and 5. informed consent to participate in the study was obtained. The exclusion criteria were: 1. diagnosis of rheumatoid arthritis, idiopathic osteonecrosis, or foot and ankle disorders; 2. individuals with bilateral toe flexion problems, neurologic diseases, or other musculoskeletal diseases that significantly impair basic movements, such as walking; and 3. Those with severe depression or dementia, which would hinder evaluation. The number of falls over one-year period was obtained retrospectively within the inclusion time by nurses. The group with one or more falls was defined as the fall group, and the group with no falls was defined as the non-fall group.

Experimental procedure

We assessed the participants one day prior to TKA. Descriptive data, fall-related assessments (presence of falls and fear of falling), radiographs, and physical function data were collected from electronic medical records. Physical function measurements were measured in the rehabilitation room, and were performed by 14 randomly assigned physical therapists for all participants to reduce bias as much as possible. The surgical side was considered as the affected side, whereas the non-surgical side was considered as the unaffected side.

Fall definition

A fall was defined as “an event that results in a person coming to rest unintentionally on the ground or other lower level, not as a result of a major intrinsic event of overwhelming hazard” [23]. Falls were excluded if they were not related to gait, standing and transfer, for example a fall with a bicycle and ladder.

Primary outcome

A toe grip dynamometer (T.K.K.3362; Takei Scientific Instruments, Niigata, Japan) was used to measure TGS (Fig 1) in a sitting position with 90° hip and knee joint flexion and the ankle in a neutral position. Under vertical loading on the foot, the plantar aponeurosis was extended with the foot truss structure [24]. The participants were instructed to place their test foot within the heel stopper and to grasp the dynamometer grip bar with their toes. They first performed a few test contractions with maximum effort to familiarize themselves with the measurement process and then performed as many voluntary isometric contractions as possible. Maximum TGS was measured twice, and the mean value (kg) was calculated. Participants performed maximum-effort contractions after the "warm-up" repetitions. An almost perfect inter- and intra-rater reliability of this measurement protocol using the toe grip dynamometer has been observed previously in people aged 60–79 years [25].

Secondary outcomes

Descriptive data, such as gender, age, height, weight, and body mass index were collected by nurses during the evaluation. The patients self-reported whether and how many times they experienced falls in the one past year. The severity of KOA was determined using the Kellgren–Lawrence (K-L) grading system [26]. Four orthopedic surgeons evaluated all medical




Fig 1. Toe grip strength assessment. a. Toe grip dynamometer employed to measure toe grip strength. b. The grip bar of the instrument was adjusted to the first metatarsophalangeal joint of the participant. The participants sat on the edge of their seats keeping their trunks in a vertical position and the hip and knee joints bent to approximately 90°.

<https://doi.org/10.1371/journal.pone.0282944.g001>

records and determined the K-L grade severity. Isometric knee extension strength (IKES) was measured using a hand-held dynamometer (μ -tas F1, ANIMA, Tokyo, Japan) with participants in a seated position and the knee in 90° flexion [27]. The reliability and validity of this measurement method have been previously demonstrated in patients with KOA [27, 28]. The participants were instructed to increase the intensity of knee extension against the dynamometer gradually and for approximately 2 seconds to avoid explosive contraction, and to maintain their maximal force output for approximately 3 seconds. Maximum IKES was measured twice, and the mean value (kg) was calculated. Pain levels at rest and during walking were determined using a visual analog scale ranging from 0 (no pain) to 100 mm (worst pain) [29]. This measurement method has been reported to be reliable and valid for the assessment of individuals with KOA [30]. The TUG was used as a behavioral measure of knee function using standard test methods [31]. The reliability and validity of this measurement method have been demonstrated [32, 33]. The participants stood up from an armless chair upon the assessor's signal, walked to the 3-m point, and returned to sit on the same chair. The TUG measurements were recorded twice and a mean value between the two measurements was calculated and used for statistical analysis. Fear of falling was assessed using the Japanese version of the modified Fall Efficacy Scale (mFES), developed by Hill et al [34]. The mFES is a 10-grade scale comprising 14 items (score range: 0–140 points), and a modified version of the Falls Efficacy Scale developed by Tinetti et al. [35]. The mFES is used to determine the level of confidence in performing specific movements and actions without falling, with higher scores indicating higher levels of self-efficacy in fall prevention and less fear of falls. The reliability and validity of this measurement method have been demonstrated [34].

Data analysis

Descriptive data were presented as the number of cases, mean with standard deviations (SD), and percentages. For all analyses, the significance level was set at 5%. All statistical analyses were performed using SPSS Statistics for Windows, version 26.0 (IBM Corp, Tokyo, Japan). The sample size was calculated by G-power Post-hoc. The sample sizes for both the fall and non-fall groups were calculated based on effect size = 0.5 and $\alpha = 0.05$.

Before comparing the difference between the two groups, a Kolmogorov–Smirnov test was performed as a homogeneity test. The results confirmed that the p-values for all factors were less than 0.05 for both groups. Therefore, Mann–Whitney- and chi-squared tests were used to test the difference between the two groups. The factors subjected to Mann–Whitney U test were age, height, weight, BMI, TGS on both sides, IKES on both sides, pain at rest on both sides, pain while walking on both sides, TUG, and mFES. Factors subjected to the chi-squared test were sex and K-L grade.

Multiple logistic regression was performed to examine the association of each factor to the dependent variable (i.e., fall or no fall history). The independent variables were height, weight, TGS on both sides, and mFES. Hosmer–Lemeshow test was performed to determine whether the analysis result was significantly fitted to the actual data. The significance of the regression equation was confirmed by rate of accurate discrimination. The variance inflation factor (VIF) was calculated to account for the degree of multicollinearity among the related factors.

Results

We excluded 77 participants, while 330 met the inclusion criteria (Fig 2). The 330 participants were divided into two groups: fall and non-fall. The participants' descriptive characteristics and results of Mann–Whitney- and chi-squared tests are summarized in Table 1. Of the 330 participants enrolled in the study, 74 (22.4%) were in the fall group and 256 (77.6%) were in the non-fall group. The percentage of fall was calculated from the 330 participants who met the inclusion criteria of the study. Of the participants with a fall history, 42 reported 1 fall, 14 reported 2 falls, 10 reported 3 falls, 4 reported 4 falls, and 4 reported 5 falls. A history of 6 or more falls was never recorded. The mean number of falls per participant was 1.85. In addition, the sample size was calculated with 74 patients in the fall group and 256 in the non-fall group, the power was 0.97.

Factors that differed significantly between the fall and non-fall groups were height ($p = .014$), TGS on the affected side ($p = .003$), TGS on the unaffected side ($p = .007$), and mFES ($p = .001$).

The results of multiple logistic regression analysis are presented in Table 2. The model chi-squared test revealed significant results, indicating association with TGS on the affected side ($\beta = -.081$, $p = .024$, Odds ratio [OR] = .922), and mFES ($\beta = -.013$, $p = .002$, OR = .987). The model χ^2 was significant at $p < .01$. The result of the Hosmer–Lemeshow test was not significant at $p = .56$, and the fit of regression equation was good. The rate of accurate discrimination was 77.6%. VIF for mFES and TGS on the affected side were 1.023 in both cases, and no multicollinearity was observed. VIF calculated from the related factors were height: 1.165; weight: 1.059; and TGS on the unaffected side: 2.320.

Discussion

We analyzed a number of factors that potentially contributed to a history of falls in patients with KOA, including TGS, and clarified which of those had an impact. Only a few studies reported the incidence of falls in individuals with KOA in Japan [36]. To the best of our knowledge, this is the first study to investigate the relationship between history of falls and TGS in older adults with KOA. Our results indicated that falls in older adults with KOA were related to TGS. In short, falls in older adults with KOA are associated with lower TGS.




Fig 2. Flow diagram of the study participants' inclusion process.

<https://doi.org/10.1371/journal.pone.0282944.g002>

The percentage of fall rate in this study

In a study of 5,062 frail older adults in Japan, approximately 30% experienced a fall at least once in a year [37]. Compared to older adults with KOA in Australia (48% fell within 12 months prior to TKA) [4] and in the United Kingdom (24% fell within 3 months prior to TKA) [5], the fall rate 12 months prior to TKA in our study was lower (22.4%). However, according to the publication of vital statistics in Japan, accidental deaths from falls among older adults are on the rise. The overall mortality rates per 100000 persons in the older population increased from 19.5 in 1997 to 20.5 in 2016 [38]. A previous study has reported that fall history before performing TKA increased the risk of post-operative falls [5, 6]. This insight might lead to a better understanding of prevention of injuries from post-operative falls.

Toe grip strength and fall history

A previous study has reported that TGS declines with age [39], resulting in diminished walking ability and static balance, which may be risk factors for falls. Tsuyuguchi et al. recruited

Table 1. Participants' descriptive characteristics and descriptive statistics for the study variables (n = 330).

	Fall group (n = 74)			Non fall group (n = 256)			p-value	Phi coefficient
Percentage of fall, %	22.4							
Gender, (female/male)	58/16			194 / 62			0.643	-0.025
Age, years	73.7	±	8.1	74.8	±	7.2	0.396	0.079
Height, cm	152.6	±	8.6	154.9	±	8.6	0.014*	
Weight, kg	61.5	±	12.9	63.0	±	12.0	0.247	
BMI, kg/m ²	26.3	±	4.0	26.2	±	3.9	0.898	
K-L grade, n (%)	II	0 (0)			5 (2.0)			0.361
	III	6 (8.1)			28 (10.9)			
	IV	68 (91.9)			223 (87.1)			
TGS, kg	affected side	7.0	±	4.3	8.6	±	4.3	0.003*
	unaffected side	7.1	±	4.0	8.7	±	4.5	0.007*
IKES, kg	affected side	15.0	±	7.2	15.8	±	9.2	0.989
	unaffected side	17.2	±	6.9	18.2	±	11.1	0.715
Rest pain, mm	affected side	15.4	±	19.6	13.1	±	20.6	0.205
	unaffected side	5.6	±	12.0	5.0	±	13.3	0.110
Walking pain, mm	affected side	50.6	±	22.3	46.2	±	26.2	0.158
	unaffected side	23.8	±	24.3	19.4	±	23.9	0.089
TUG, s	12.6	±	4.2	12.8	±	4.5	0.989	
mFES, point	98.4	±	35.5	113.0	±	28.1	0.001*	

Data were expressed as mean (standard deviation). *P<0.05

BMI, Body mass index; K-L grade, Kellgren-Lawrence grade; TGS, Toe grip strength; IKES, Isometric knee extension strength; TUG, Timed up and go test; mFES, modified Fall Efficacy Scale

<https://doi.org/10.1371/journal.pone.0282944.t001>

middle-aged adults and, whose average age was 62.02, divide them into high and low risk of falls. They found TGS to be an independent risk factor for fall occurrence [40]. However, there is no report on whether reduced TGS in individuals with KOA is associated with the fall itself. Based on the results of our study, we believe that TGS contributes to the challenges faced by older adults with KOA and a fall history. Although a detailed causal relationship is unknown, multiple regression analysis has identified TGS as an independent factor associated with KOA [41]. Conversely, abnormal loading of the knee joint can be caused by changes in the kinematic relationship between the foot and knee [42, 43]. Compared to healthy older adults, those with KOA have lower TGS, and the measured pressure decreases during walking [44, 45]. It is possible that the progression of KOA leads to decreased TGS; conversely, decreased TGS may contribute to KOA progression. However, this causal relationship is unclear; therefore, further studies are required to investigate it. Regardless of the causal pathway, there is interdependence between TGS and KOA, which increases the risk of falls. In the future, studies should

Table 2. Results of the multiple logistic regression analysis.

	B	Standard Error	Wald	p-value	Odds ratios (95% confidence interval)
TGS on the affected side	-0.081	0.036	5.069	0.024	0.922 (0.86–0.99)
mFES	-0.013	0.004	9.673	0.002	0.987 (0.98–0.99)
Constant	0.760	0.494	2.370	0.124	

Regression equation: $\ln(p/1-p) = \{-0.081(\text{TGS on the affected side})\} + \{-0.013(\text{mFES})\} + 0.760$ model χ^2 , $p < .01$; Hosmer-Lemeshow test, $P = .56$; rate of accurate discrimination, 77.6% TGS, Toe grip strength; mFES, modified Fall Efficacy Scale

<https://doi.org/10.1371/journal.pone.0282944.t002>

approach the causal relationship between KOA and TGS from the perspectives of kinesiology and biomechanics, in addition to seeking strategies to prevent falls.

Self-efficacy for falls and fall history

Adults with KOA with more frequent falls may have a more pronounced fear of falling than those with fewer falls. Tinetti et al. [46] defined fear of falling as "anxiety about falling that causes one to avoid activities of daily living, even though one is capable of performing them". In their study, fear of falling depended on the history of falls, ranging from 12–65% among community-dwelling older adults without a fall history, and from 29–92% among those with [47]. Therefore, fall history is associated with fear of falling [48]. Our results support those of previous studies [48]. In addition, participants with KOA and a history of falls experienced knee pain, knee instability, and muscle weakness in the lower extremities, which could have also contributed to a greater fear of falling. Post-fall syndrome, in which a loss of self-confidence after a fall leads to a decreased activity level, in turn amplifies the fear of falling.

Limitations

This study had three major limitations. First, TGS was the only assessment performed on the foot. The degree of flatfoot and range of motion of the foot, which are common problems in individuals with KOA, were not measured. The abnormal foot posture of KOA has excessive first medial tibiofemoral contact force during walking [49]. As these factors are also associated with gait, which is involved in half of all fall scenarios [50], they are likely to contribute to falls. Second, previous studies have reported decreased physical activity [51] as well as hip [52] and ankle [53] weakness as risk factors for falls. Because we did not evaluate these factors and did not include these results in our regression analysis, we could not determine the relative contribution of TGS to physical activity and muscle strength for the above factors in older adults with KOA. Finally, we conducted a cross-sectional observational study; therefore, subsequent cohort and intervention studies should be conducted to better clarify the causal relationships between foot function and falls in older adults with KOA.

Conclusions

In this study, we investigated the factors leading to falls in older adults with KOA in Japan. Our results indicated that falls in older adults with KOA were related to TGS. In short, falls in older adults with KOA are associated with lower TGS. Physical therapy interventions to enhance TGS could be one solution to help prevent falls in individuals with KOA. In the future, cohort and interventional studies evaluating the relationship between TGS and falls should be performed.

Supporting information

S1 Checklist. STROBE statement—checklist of items that should be included in reports of cross-sectional studies.

(DOCX)

S1 Dataset.

(XLSX)

Acknowledgments

We would like to thank Editage (www.editage.jp) for English language editing.

Author Contributions

Conceptualization: Yuya Mawarikado, Yusuke Inagaki, Akira Kido, Yasuhito Tanaka.

Data curation: Yuya Mawarikado, Yusuke Inagaki.

Formal analysis: Yuya Mawarikado, Yusuke Inagaki.

Investigation: Yuya Mawarikado, Yusuke Inagaki, Takanari Kubo.

Methodology: Yuya Mawarikado, Yusuke Inagaki, Takanari Kubo.

Project administration: Yuya Mawarikado.

Resources: Yuya Mawarikado, Tadashi Fujii.

Software: Yuya Mawarikado, Yusuke Inagaki.

Visualization: Yuya Mawarikado.

Writing – original draft: Yuya Mawarikado, Yusuke Inagaki, Akira Kido.

Writing – review & editing: Yasuhito Tanaka.

References

1. Gillespie LD, Gillespie WJ, Robertson MC, Lamb SE, Cumming RG, Rowe BH. Interventions for preventing falls in elderly people. *Cochrane Database Syst Rev*. 2003; 4: CD000340. <https://doi.org/10.1002/14651858.CD000340> PMID: 14583918
2. Doré AL, Golightly YM, Mercer VS, Shi XA, Renner JB, Jordan JM, et al. Lower-extremity osteoarthritis and the risk of falls in a community-based longitudinal study of adults with and without osteoarthritis. *Arthritis Care Res (Hoboken)*. 2015; 67: 633–639. <https://doi.org/10.1002/acr.22499> PMID: 25331686
3. Williams SB, Brand CA, Hill KD, Hunt SB, Moran H. Feasibility and outcomes of a home-based exercise program on improving balance and gait stability in women with lower-limb osteoarthritis or rheumatoid arthritis: A pilot study. *Arch Phys Med Rehabil*. 2010; 91: 106–114. <https://doi.org/10.1016/j.apmr.2009.08.150> PMID: 20103404
4. Levinger P, Menz HB, Wee E, Feller JA, Bartlett JR, Bergman NR. Physiological risk factors for falls in people with knee osteoarthritis before and early after knee replacement surgery. *Knee Surg Sports Traumatol Arthrosc*. 2011; 19: 1082–1089. <https://doi.org/10.1007/s00167-010-1325-8> PMID: 21107530
5. Swinkels A, Newman JH, Allain TJ. A prospective observational study of falling before and after knee replacement surgery. *Age Ageing*. 2009; 38: 175–181. <https://doi.org/10.1093/ageing/afn229> PMID: 19029071
6. Chan ACM, Jehu DA, Pang MYC. Falls after total knee arthroplasty: frequency, circumstances, and associated factors—a prospective cohort study. *Phys Ther*. 2018; 98: 767–778. <https://doi.org/10.1093/ptj/pzy071> PMID: 29931164
7. Manlapaz DG, Sole G, Jayakaran P, Chapple CM. Risk factors for falls in adults with knee osteoarthritis: a systematic review. *PM R*. 2019; 11: 745–757. <https://doi.org/10.1002/pmrj.12066> PMID: 30609282
8. Aljehani MS, Crenshaw JR, Rubano JJ, Dellose SM, Zeni JA Jr. Falling risk in patients with end-stage knee osteoarthritis. *Clin Rheumatol*. 2021; 40: 3–9. <https://doi.org/10.1007/s10067-020-05165-6> PMID: 32500227
9. Herman T, Giladi N, Hausdorff JM. Properties of the 'Timed UP and Go' test: more than meets the eye. *Gerontology*. 2011; 57: 203–210. <https://doi.org/10.1159/000314963> PMID: 20484884
10. Kang L, Han P, Wang J, Ma Y, Jia L, Fu L, et al. Timed UP and Go test can predict recurrent falls: a longitudinal study of the community-dwelling elderly in China. *Clin Interv Aging*. 2017; 12: 2009–2016. <https://doi.org/10.2147/CIA.S138287> PMID: 29238175
11. Loft CC, Jones FW, Kneebone II. Falls self-efficacy and falls incidence in community-dwelling older people: the mediating role of coping. *Int Psychogeriatr*. 2018; 30: 727–733. <https://doi.org/10.1017/S1041610217002319> PMID: 29113615
12. Ahmadiyahangar A, Javadian Y, Babaei M, Heidari B, Hosseini S, Aminzadeh M. The role of quadriceps muscle strength in the development of falls in the elderly people, a cross-sectional study. *Chiropr Man Therap*. 2018; 26: 31. <https://doi.org/10.1186/s12998-018-0195-x> PMID: 30128111

13. Menz HB, Morris ME, Lord SR. Foot and ankle characteristics associated with impaired balance and functional ability in older people. *J Gerontol A Biol Sci Med Sci*. 2005; 60: 1546–1552. <https://doi.org/10.1093/gerona/60.12.1546> PMID: 16424286
14. Menz HB, Morris ME, Lord SR. Foot and ankle risk factors for falls in older people: a prospective study. *J Gerontol A Biol Sci Med Sci*. 2006; 61: 866–870. <https://doi.org/10.1093/gerona/61.8.866> PMID: 16912106
15. Mickle KJ, Munro BJ, Lord SR, Menz HB, Steele JR. ISB Clinical Biomechanics Award 2009: toe weakness and deformity increase the risk of falls in older people. *Clin Biomech (Bristol, Avon)*. 2009; 24: 787–791. <https://doi.org/10.1016/j.clinbiomech.2009.08.011> PMID: 19751956
16. Endo M, Ashton-Miller JA, Alexander NB. Effects of age and gender on toe flexor muscle strength. *J Gerontol A Biol Sci Med Sci*. 2002; 57: M392–M397. <https://doi.org/10.1093/gerona/57.6.m392> PMID: 12023269
17. Menz HB, Zammit GV, Munteanu SE, Scott G. Plantarflexion strength of the toes: age and gender differences and evaluation of a clinical screening test. *Foot Ankle Int*. 2006; 27: 1103–1108. <https://doi.org/10.1177/107110070602701217> PMID: 17207439
18. Scott G, Menz HB, Newcombe L. Age-related differences in foot structure and function. *Gait Posture*. 2007; 26: 68–75. <https://doi.org/10.1016/j.gaitpost.2006.07.009> PMID: 16945538
19. Misu S, Doi T, Asai T, Sawa R, Tsutsumimoto K, Nakakubo S, et al. Association between toe flexor strength and spatiotemporal gait parameters in community-dwelling older people. *J Neuroeng Rehabil*. 2014; 11: 143. <https://doi.org/10.1186/1743-0003-11-143> PMID: 25297707
20. Cebolla EC, Rodacki AL, Bento PC. Balance, gait, functionality and strength: comparison between elderly fallers and non-fallers. *Braz J Phys Ther*. 2015; 19: 146–151. <https://doi.org/10.1590/bjpt-rbf.2014.0085> PMID: 25993628
21. Kataoka H, Miyatake N, Kitayama N, Murao S, Kohi F, Tanaka S. Relationship of toe pinch force to other muscle strength parameters in men with type 2 diabetes. *Environ Health Prev Med*. 2016; 21: 179–185. <https://doi.org/10.1007/s12199-016-0512-8> PMID: 26879315
22. Tsuyuguchi R, Kurose S, Seto T, Takao N, Fujii A, Tsutsumi H, et al. The effects of toe grip training on physical performance and cognitive function of nursing home residents. *J Physiol Anthropol*. 2019; 38: 11. <https://doi.org/10.1186/s40101-019-0202-5> PMID: 31462272
23. Clark RD, Lord SR, Webster IW. Clinical parameters associated with falls in an elderly population. *Gerontology*. 1993; 39(2):117–123. <https://doi.org/10.1159/000213521> PMID: 8514201
24. Sarrafian SK. Functional characteristics of the foot and plantar aponeurosis under tibiotalar loading. *Foot Ankle*. 1987; 8: 4–18. <https://doi.org/10.1177/107110078700800103> PMID: 3623360
25. Uritani D, Fukumoto T, Matsumoto D. Intrarater and interrater reliabilities for a toe grip dynamometer. *J Phys Ther Sci*. 2012; 24: 639–643. <https://doi.org/10.1589/jpts.24.639>
26. Kellgren JH, Lawrence JS. Radiological assessment of osteo-arthrosis. *Ann Rheum Dis*. 1957; 16: 494–502. <https://doi.org/10.1136/ard.16.4.494> PMID: 13498604
27. Koblbauer IF, Lambrecht Y, Van der Hulst ML, Neeter C, Engelbert RH, Poolman RW, et al. Reliability of maximal isometric knee strength testing with modified hand-held dynamometry in patients awaiting total knee arthroplasty: useful in research and individual patient settings? A reliability study. *BMC Musculoskelet Disord*. 2011; 12: 249. <https://doi.org/10.1186/1471-2474-12-249> PMID: 22040119
28. Chopp-Hurley JN, Wiebenga EG, Gatti AA, Maly MR. Investigating the test-retest reliability and validity of hand-held dynamometry for measuring knee strength in older women with knee osteoarthritis. *Physiother Can*. 2019; 71: 231–238. <https://doi.org/10.3138/ptc-2018-0051> PMID: 31719719
29. McCormack HM, Horne DJ, Sheather S. Clinical applications of visual analogue scales: a critical review. *Psychol Med*. 1988; 18: 1007–1019. <https://doi.org/10.1017/s0033291700009934> PMID: 3078045
30. Alghadir AH, Anwer S, Iqbal A, Iqbal ZA. Test-retest reliability, validity, and minimum detectable change of visual analog, numerical rating, and verbal rating scales for measurement of osteoarthritic knee pain. *J Pain Res*. 2018; 11: 851–856. <https://doi.org/10.2147/JPR.S158847> PMID: 29731662
31. Podsiadlo D, Richardson S. The Timed “UP & Go”: a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc*. 1991; 39: 142–148. <https://doi.org/10.1111/j.1532-5415.1991.tb01616.x> PMID: 1991946
32. Alghadir A, Anwer S, Brismée JM. The reliability and minimal detectable change of Timed UP and Go test in individuals with grade 1–3 knee osteoarthritis. *BMC Musculoskelet Disord*. 2015; 16: 174. <https://doi.org/10.1186/s12891-015-0637-8> PMID: 26223312
33. Nightingale CJ, Mitchell SN, Butterfield SA. Validation of the Timed UP and Go Test for assessing balance variables in adults aged 65 and older. *J Aging Phys Act*. 2019; 27: 230–233. <https://doi.org/10.1123/japa.2018-0049> PMID: 30117359

34. Hill KD, Schwarz JA, Kalogeropoulos AJ, Gibson SJ. Fear of falling revisited. *Arch Phys Med Rehabil.* 1996; 77: 1025–1029. [https://doi.org/10.1016/s0003-9993\(96\)90063-5](https://doi.org/10.1016/s0003-9993(96)90063-5) PMID: 8857881
35. Tinetti ME, Richman D, Powell L. Falls efficacy as a measure of fear of falling. *J Gerontol.* 1990; 45: P239–P243. <https://doi.org/10.1093/geronj/45.6.p239> PMID: 2229948
36. Muraki S, Akune T, Ishimoto Y, Nagata K, Yoshida M, Tanaka S, et al. Risk factors for falls in a longitudinal population-based cohort study of Japanese men and women: the ROAD Study. *Bone.* 2013; 52: 516–523. <https://doi.org/10.1016/j.bone.2012.10.020> PMID: 23103329
37. Shimada H, Suzukawa M, Ishizaki T, Kobayashi K, Kim H, Suzuki T. Relationship between subjective fall risk assessment and falls and fall-related fractures in frail elderly people. *BMC Geriatr.* 2011; 11: 40. <https://doi.org/10.1186/1471-2318-11-40> PMID: 21838891
38. Hagiya H, Koyama T, Zamami Y, Tatebe Y, Funahashi T, Shinomiya K, et al. Fall-related mortality trends in older Japanese adults aged ≥ 65 years: a nationwide observational study. *BMJ, (Open).* 2019; 9: e033462. <https://doi.org/10.1136/bmjopen-2019-033462> PMID: 31831549
39. Uritani D, Fukumoto T, Matsumoto D, Shima M. Reference values for toe grip strength among Japanese adults aged 20 to 79 years: a cross-sectional study. *J Foot Ankle Res.* 2014; 7: 28. <https://doi.org/10.1186/1757-1146-7-28> PMID: 24855492
40. Tsuyuguchi R, Kurose S, Seto T, Takao N, Tagashira S, Tsutsumi H, et al. Toe grip strength in middle-aged individuals as a risk factor for falls. *J Sports Med Phys Fitness.* 2018; 58: 1325–1330. <https://doi.org/10.23736/S0022-4707.17.07473-4> PMID: 28738667
41. Uritani D, Fukumoto T, Myodo T, Fujikawa K, Usui M, Tataro D. The association between toe grip strength and osteoarthritis of the knee in Japanese women: a multicenter cross-sectional study. *PLOS ONE.* 2017; 12: e0186454. <https://doi.org/10.1371/journal.pone.0186454> PMID: 29040330
42. Lafortune MA, Cavanagh PR, Sommer HJ 3rd, Kalenak A. Foot inversion-eversion and knee kinematics during walking. *J Orthop Res.* 1994; 12: 412–420. <https://doi.org/10.1002/jor.1100120314> PMID: 8207595
43. Souza TR, Pinto RZ, Trede RG, Kirkwood RN, Fonseca ST. Temporal couplings between rearfoot-shank complex and hip joint during walking. *Clin Biomech (Bristol, Avon).* 2010; 25: 745–748. <https://doi.org/10.1016/j.clinbiomech.2010.04.012> PMID: 20621756
44. Saito I, Okada K, Nishi T, Wakasa M, Saito A, Sugawara K, et al. Foot pressure pattern and its correlation with knee range of motion limitations for individuals with medial knee osteoarthritis. *Arch Phys Med Rehabil.* 2013; 94: 2502–2508. <https://doi.org/10.1016/j.apmr.2013.07.017> PMID: 23916619
45. Arnold JB, Mackintosh S, Jones S, Thewlis D. Differences in foot kinematics between young and older adults during walking. *Gait Posture.* 2014; 39: 689–694. <https://doi.org/10.1016/j.gaitpost.2013.09.021> PMID: 24183676
46. Tinetti ME, Powell L. Fear of falling and low self-efficacy: a case of dependence in elderly persons. *J Gerontol.* 1993; 48;Spec No: 35–38. https://doi.org/10.1093/geronj/48.special_issue.35 PMID: 8409238
47. Legters K. Fear of falling. *Phys Ther.* 2002; 82: 264–272. <https://doi.org/10.1093/ptj/82.3.264> PMID: 11869155.
48. Cumming RG, Salkeld G, Thomas M, Szonyi G. Prospective study of the impact of fear of falling on activities of daily living, SF-36 scores, and nursing home admission. *J Gerontol A Biol Sci Med Sci.* 2000; 55: M299–M305. <https://doi.org/10.1093/gerona/55.5.m299> PMID: 10819321
49. Kubo T, Uritani D, Ogaya S, Kita S, Fukumoto T, Fujii T, et al. Association between foot posture and tibiofemoral contact forces during barefoot walking in patients with knee osteoarthritis. *BMC Musculoskelet Disord.* 2022; 23: 660. <https://doi.org/10.1186/s12891-022-05624-y> PMID: 35820878
50. Li W, Keegan TH, Sternfeld B, Sidney S, Quesenberry CP, Kelsey JL. Outdoor falls among middle-aged and older adults: a neglected public health problem. *Am J Public Health.* 2006; 96: 1192–1200. <https://doi.org/10.2105/AJPH.2005.083055> PMID: 16735616
51. Di Pietro L, Campbell WW, Buchner DM, Erickson KI, Powell KE, Bloodgood B, et al. Physical Activity, injurious falls, and physical function in aging: an umbrella review. *Med Sci Sports Exerc.* 2019; 51: 1303–1313. <https://doi.org/10.1249/MSS.0000000000001942> PMID: 31095087
52. Arvin M, van Dieën JH, Faber GS, Pijnappels M, Hoozemans MJM, Verschueren SMP. Hip abductor neuromuscular capacity: a limiting factor in mediolateral balance control in older adults? *Clin Biomech (Bristol Avon).* 2016; 37: 27–33. <https://doi.org/10.1016/j.clinbiomech.2016.05.015> PMID: 27286555
53. Spink MJ, Fotoohabadi MR, Wee E, Hill KD, Lord SR, Menz HB. Foot and ankle strength, range of motion, posture, and deformity are associated with balance and functional ability in older adults. *Arch Phys Med Rehabil.* 2011; 92: 68–75. <https://doi.org/10.1016/j.apmr.2010.09.024> PMID: 21187207